



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Endovascular treatment of complex aortic injuries 40 years after surgery for aortic coarctation

Caporali, Elena ; Van Den Berg, Jos C ; Ferrari, Enrico

DOI: <https://doi.org/10.1016/j.jvscit.2018.02.013>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-158789>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Caporali, Elena; Van Den Berg, Jos C; Ferrari, Enrico (2018). Endovascular treatment of complex aortic injuries 40 years after surgery for aortic coarctation. *Journal of Vascular Surgery Cases and Innovative Techniques*, 4(2):181.

DOI: <https://doi.org/10.1016/j.jvscit.2018.02.013>


Endovascular treatment of complex aortic injuries

40 years after surgery for aortic coarctation

Elena Caporali, MD,^a Jos C. Van Den Berg, MD,^b and Enrico Ferrari, MD,^a Lugano, Switzerland

We describe the case of a 57-year-old woman suffering from systemic hypertension and thoracic pain who was operated on for aortic coarctation at the age of 16 years. Computed tomography angiography showed a pseudoaneurysm of the thoracic aorta at the level of the distal anastomosis of the previously implanted Dacron vascular graft, a penetrating ulcer of the descending thoracic aorta (distal from the graft), and an aneurysm of the left subclavian artery presumably at the level of the vascular clamp position during previous surgery (A). A custom-made endoprosthesis (Bolton Medical Products, Shrewsbury, Mass) with a proximal diameter of 22 mm, waist diameter of 18 mm, and distal diameter of 30 mm and a covered length of 170 mm was designed to perfectly fit within the previously implanted surgical Dacron vascular graft of 16 mm in diameter. The patient underwent surgical left carotid-subclavian bypass grafting with reimplantation of the left vertebral artery during the time needed to manufacture the custom-made device. Subsequently, the endovascular procedure was performed in a hybrid operating room with proper positioning (B) and release (C) of the custom-made endoprosthesis. The postoperative course was uneventful with discharge home after 5 days. Computed tomography angiography performed after 1 month confirmed a good result and exclusion of all aneurysms (D).

Submitted Feb 8, 2018; accepted Feb 24, 2018.



From the Cardiac Surgery Department, Cardiocentro Ticino Foundation^a, and the Interventional Radiology Unit, Lugano City Hospital.^b

Author conflict of interest: none.

E-mail: elena.caporali@cardiocentro.org.

The editors and reviewers of this article have no relevant financial relationships to disclose per the Journal policy that requires reviewers to decline review of any manuscript for which they may have a conflict of interest.

J Vasc Surg Cases and Innovative Techniques 2018;4:181

2468-4287

© 2018 Published by Elsevier Inc. on behalf of Society for Vascular Surgery. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<https://doi.org/10.1016/j.jvscit.2018.02.013>